

Uncertainty in Satellite estimate of Global Mean Sea Level changes, trend and acceleration

Michaël Ablain^{1,3}, Benoît Meyssignac², Lionel Zawadzki³, Rémi Jugier¹, Aurélien Ribes⁴, Giorgio Spada⁵, Jérôme Benveniste⁶, Anny Cazenave², Nicolas Picot⁷

5 ¹ MAGELLIUM, Ramonville Saint-Agne, 31520, France

² LEGOS, CNES, CNRS, IRD, Université Paul Sabatier, Toulouse, France

³ Collecte Localisation Satellite (CLS), Ramonville Saint-Agne, 31520, France

⁴ CNRM, Université Paul Sabatier, Météo France, CNRS, Toulouse, France

⁵ Dipartimento di Scienze Pure e Applicate (DiSPeA), Urbino University "Carlo Bo", 61029, Urbino, PU, Italy

10 ⁶ European Space Agency (ESA-ESRIN), Frascati, Italy

⁷ Centre National d'Études Spatiales (CNES), Toulouse, 31400, France

Correspondence to: Michaël Ablain (michael.ablain@magellium.fr)

Abstract. Satellite altimetry missions now provide more than 25 years of accurate, continuous and quasi-global
15 measurements of sea level along the reference ground track of TOPEX/Poseidon. These measurements are used
by different groups to build the Global Mean Sea Level (GMSL) record, an essential climate change indicator.
Estimating a realistic uncertainty of the GMSL record is of crucial importance for climate studies such as assessing
precisely the current rate and acceleration of sea level, analysing the closure of the sea level budget, understanding
the causes of sea level rise, detecting and attributing the response of sea level to anthropogenic activity, or
20 calculating the Earth's energy imbalance. Previous authors have estimated the uncertainty of the GMSL trend over
the period 1993-2014 by thoroughly analysing the error budget of the satellite altimeters and have shown that it
amounts to ± 0.5 mm/yr (90% confidence level). In this study, we extend our previous results providing a
comprehensive description of the uncertainties in the satellite GMSL record. We analysed 25 years of satellite
25 altimetry data and provided for the first time the error variance-covariance matrix for the GMSL record with a time
resolution of ten days. Three types of errors have been modelled (drifts, biases, noises) and combined together to
derive a realistic estimate of the GMSL error variance-covariance matrix. From the latter, we derived a 90%
confidence envelop of the GMSL record on a 10-day basis. Then we used a least squared approach and the error
variance-covariance matrix to assess the GMSL trend and acceleration uncertainties over any 5-year time periods
and longer in between October 1992 and December 2017. Over 1993-2017, we have found a GMSL trend of
30 3.35 ± 0.4 mm/yr within a 90% Confidence Level (CL) and a GMSL acceleration of 0.12 ± 0.07 mm/yr² (90% CL).
This is in agreement (within error bars) with previous studies. The full GMSL error variance-covariance matrix is
freely available online: <https://doi.org/10.17882/58344> (Ablain et al., 2018).

1 Introduction

The sea level change is a key indicator of global climate change, which integrates changes in several components of the climatic system as a response to climatic variability, both anthropogenic and natural. Since October 1992, sea level variations have been routinely measured by twelve high-precision altimeter satellites providing more than 25 years of continuous measurements. The altimeter Global Mean Sea Level (GMSL) indicator is calculated from the accurate and stable measurements of four reference altimeter missions, namely TOPEX/Poseidon (T/P), Jason-1, Jason-2 and Jason-3. All four reference missions are flying (or have flown) over the same historical ground track on a 10-day repeat cycle. They all have been precisely inter-calibrated (Zawadzki and Ablain, 2016) to ensure the long term stability of the sea level measurements. Six research groups (AVISO/CNES, SL_cci/ESA, University of Colorado, CSIRO, NASA/GSFC, NOAA) have processed the sea level raw data provided by satellite altimetry to provide the GMSL series on a 10-day basis (Figure 1). The six different estimates of the GMSL record show small deviations between 1 and 2 mm at inter-annual time scales (1 to 5-year time scales) and between ± 0.15 mm/yr in terms of trend over the period 1993-2017. The spread across these estimates is due to the use of various processing techniques, alternative versions of ancillary data and different interpolation methods applied by the several groups (Masters et al., 2012, Henry et al., 2014). This spread is smaller than the real uncertainty in the sea level trend, because all the research groups have used similar methods and corrections to process the raw data and thus several sources of systematic uncertainty are not accounted for in the spread.

In a previous study Ablain et al. (2009) have proposed a realistic estimate of the uncertainty in the GMSL trend over 1993-2008, using an approach based on the error budget. They have identified the radiometer wet tropospheric correction as one of the main sources of error. They have also proposed the orbital determination, the inter-calibration of altimeters and the estimate of the altimeter range, σ_0 and significant wave height (mainly on TOPEX/Poseidon) as significant sources of error. When all the terms were accounted for, they have found that the uncertainty on the trend over 1993-2008 was ± 0.6 mm/yr within a 90% confidence level. This is larger than the uncertainty of ± 0.3 mm/yr over a 10-year period required by GCOS (GCOS, 2011). In the framework of the ESA Sea Level Climate Change Initiative (SL_cci), significant improvements have been obtained estimating the sea level from space (Ablain et al., 2015; Quartly et al., 2017, Legeais et al., 2018) to get closer to the GCOS requirements. New altimeter standards including new wet troposphere corrections, new orbit solutions, new atmospheric corrections and others, were selected and applied in order to improve the sea level estimation. The GMSL trend uncertainties were then updated and estimated at different temporal and spatial scales (Ablain et al., 2015; Legeais et al., 2018). During the second altimetry decade (2002- 2014), Ablain et al. (2015) have estimated that the uncertainty of the GMSL trend was lower than ± 0.5 mm/yr within a 90% Confidence Level (CL) for periods longer than 10 years.

In previous studies, the uncertainty in GMSL has been assessed for long-term trends (periods of 10 years or more, starting in 1993), inter-annual time scales (between 1 and 5 years) and annual time scales (Ablain et al., 2009; 2015). This estimation of the uncertainty at three time-scales is a valuable first step, but it is not enough, as it does not fully meet the needs of the scientific community. In many climatic studies the GMSL uncertainty is required at different time scales and span within the 25-year altimetry record. In sea level budget studies based on the evolution of GMSL components, these estimates have been carried out at monthly time scale. In this way, the GMSL monthly changes have been interpreted in terms of changes of ocean mass (Gravity recovery and climate experiment – GRACE – mission). This is also the case of studies estimating the Earth's energy imbalance with the sea-level budget approach (Meysignac et al., 2018). In the studies on the detection and the attribution of climate change (e.g. Slangen et al., 2017), the uncertainty in the trend estimates is needed, but over different time spans that those addressed in Ablain et al. (2015, 2009) and in Legeais et al. (2018). The uncertainty on different metrics is often needed. Dieng et al. (2017) and Nerem et al. (2018) have recently estimated the acceleration in the GMSL over 1993-2017, finding a small acceleration (0.08 mm/yr^2) over the 25-year long altimetry record.

80

In this paper we focus on the uncertainty in the GMSL record arising from instrumental errors in the satellite altimetry. The uncertainties of the measurements have been quantified in the GMSL record. This is an important information for the studies in detection and attribution of the climatic changes, estimating the GMSL rise as a response to the anthropogenic activity. But this is not sufficient. In the detection-attribution studies the response of the GMSL to the anthropogenic activity needs to be separated from the response to the natural variability of the climate system, representing an additional source of uncertainty.

The objective of this paper is to estimate the error variance-covariance matrix of the GMSL (on a 10-day basis) from satellite altimetry measurements. This error variance-covariance matrix provides a comprehensive description of the uncertainties in GMSL to users. It covers all time scales that are included in the 25-year long satellite altimetry record: from 10 days (the time resolution of the GMSL time series) to multidecadal time scales. It also enables us to estimate the uncertainty in any metric derived from GMSL measurements such as trend, acceleration or other moments of higher order in a consistent way.

We used an error budget approach to a global scale on a 10-day basis in order to calculate the error variance-covariance matrix. We considered all the major sources of uncertainty in the altimetry measurements, including the wet tropospheric correction, the orbital solutions, the inter-calibration of satellites. We have also taken into account the time correlation between the different sources of uncertainty (section 2). The errors have been separately characterized for each altimetry mission, since they have been affected by different sources of uncertainty (section 2). On the basis of the error variance-covariance matrix we estimate the uncertainty in GMSL individual

100 measurements on a 10-day basis (section 3) and the uncertainty in trend and acceleration over all periods included
in the 25-year satellite altimetry record (1993-2017) (Section 4). Note that in this article all uncertainties associated
to the GMSL are reported with a 90% CL unless stated otherwise.

1 GMSL data series

The six main groups that provide satellite altimetry based GMSL estimates (AVISO/CNES, SL_cci/ESA, University
105 of Colorado, CSIRO, NASA/GSFC, NOAA) use 1-Hz altimetry measurements from T/P, Jason-1, Jason-2 and
Jason-3 missions from 1993 to 2018 (1993-2015 for SL_cci/ESA). Each group processes the 1-Hz data with
geophysical corrections to correct the altimetry measurements for various aliasing, biases and drifts caused by
different atmospheric condition, sea states, ocean tides and others (Ablain et al., 2009). They spatially average the
data over each 10-day orbital cycle to provide GMSL time series on a 10-day basis. The differences among the
110 GMSL estimates from several groups arise from data editing, from differences in the geophysical corrections and
from differences in the used method to spatially average individual measurements during the orbital cycles (
Masters et al., 2012; Henry et al., 2014;).

Recently, the comparisons of the GMSL time series derived from satellite altimetry with independent estimates are
based on tide gauge records (Valladeau et al., 2012; Watson et al., 2015) or on the combination of the contribution
115 to sea level from thermal expansion, land ice melt and land water storage (Dieng et al., 2017). They have shown
that there was a drift in the GMSL record over the period 1993-1998. This drift is caused by an erroneous on-board
calibration correction on TOPEX altimeter side-A (noted TOPEX-A). TOPEX-A was operated from launch in
October 1992 to the end of January 1999. Then TOPEX side-B altimeter (noted TOPEX-B) took over in February
1999 (Beckley et al., 2017). The impact on the GMSL changes is -1.0 mm/yr between January 1993 and July 1995,
120 and +3.0 mm/yr between August 1995 and February 1999, with an uncertainty of ± 1.7 mm/yr (within a 90%CL,
(Ablain, 2017)).

Without taking into account the TOPEX-A drift correction, the differences between all GMSL time series are small.
The maximum trend difference between all-time series over 1993-2017 is lower than 0.15 mm/yr, representing less
125 than 5% of the GMSL trend. The differences observed at interannual time scales are also small (<2 mm). By
correcting the drift of TOPEX-A using either of the available empirical corrections (WCRP Global Sea Level Budget
Group, 2018) the differences among solutions remain the same (the difference between empirical corrections being
smaller than the difference between the raw GMSL time series).. Therefore, the choice of one or the other GMSL
record is not decisive in this study, whose purpose is to characterize the uncertainties. Hereafter, we use the GMSL
130 AVISO record. The corresponding altimeter standard corrections and the GMSL processing methods are described
on the AVISO website (<https://www.aviso.altimetry.fr/msl/>).

2 Altimetry GMSL error budget

135 This section describes the different errors that affect the altimetry GMSL record. It builds on the GMSL error budget
presented in Ablain et al. (2009) and extends this work by taking into account the new altimeter missions (Jason-
2, Jason-3) and the recent findings on altimetry error estimates. Three types of errors are considered: a) biases in
GMSL between successive altimetry missions which are characterized by bias uncertainties ($\pm\Delta$) at a given time
(t); b) drifts in GMSL characterized by a trend uncertainty ($\pm\delta$) and c) other measurement errors which exhibit time-
140 correlation (so called residual time correlated errors here after). The residual time correlated errors are
characterized by their standard deviation (σ) and by the correlation time-scale (λ). All altimetry errors identified in
this study are summarized in **Erreur ! Source du renvoi introuvable.** and detailed hereafter. Note that all
uncertainties reported in Table 1 are Gaussians and they are given at the 1-sigma level (i.e. we provide the standard
deviation of the Gaussian, noted $1-\sigma$ hereafter).

145 The biases can arise between the GMSL record of two successive satellite missions like between T/P and Jason-
1 in May 2002, Jason-1 and Jason-2 in October 2008 and between Jason-2 and Jason-3 in October 2016. These
biases are estimated during dedicated 9-month inter-calibration phases when a satellite altimeter and its successor
fly over the same track, 1 minute apart. During the inter-calibration phases the bias is estimated and corrected for.
Different missions show different biases, but the uncertainty in the bias correction is the same for all inter-calibration
150 phases and amounts: ± 0.5 mm (Zawadzki and Ablain, 2016). The situation is different for the switch from TOPEX-
A to TOPEX-B in February 1999 because it was impossible to do any inter-calibration phase between the two sides
of TOPEX (as both instruments were flying on the same spacecraft). For the switch, we assume that the uncertainty
in GMSL is larger and is about 2 mm (Zawadzki and Ablain, 2016).

155 The drifts may occur in the GMSL record because of drifts in TOPEX-A and TOPEX-B radar instruments, because
of drifts in the International Terrestrial Reference Frame (ITRF) realization in which altimeter orbits are determined
or because of drifts in the Glacial Isostatic Adjustment (GIA) correction applied to the GMSL record. As explained
before, the TOPEX-A record shows a spurious drift due to an erroneous on-board calibration correction of the
altimeter (Beckley et al., 2017). This drift has been corrected by using several empirical approaches (Ablain, 2017;
160 Beckley et al., 2017; Dieng et al., 2017), that are all affected by a significant uncertainty. We estimated this
uncertainty to be ± 0.7 mm/yr ($1-\sigma$ level) over the TOPEX-A period (1993-1998), with a comparison against an
independent GMSL estimate based on tide gauge records (Ablain, 2017). For the TOPEX-B record, no GMSL drift
has been reported, but Ablain et al. (2012) showed significant Sigma-0 instabilities in the order of 0.1 dB, which
generate through the sea-state bias correction an uncertainty of ± 0.1 mm/yr ($1-\sigma$ level) in the GMSL record over

165 the TOPEX-B period (February 1999 – April 2002). Concerning the ITRF realization, Couhert et al. (2015) have shown that the uncertainty on the ITRF realization drift generates an uncertainty of ± 0.1 mm/yr ($1-\sigma$ level) on the GMSL trend over 1993-2015. We adopt this value here for the whole period 1993-2017. For the uncertainty on the GIA correction applied to the GMSL, we use the value of 0.05 mm/yr ($1-\sigma$ level) over the altimetry period from Spada (2017) (the value is taken from the table 1 in Spada (2017)). It has been confirmed recently with an ensemble
170 of 1000 GIA runs, see [Melini and Spada, 2019](#)). Combining the uncertainty on the GMSL trend over 1993-2017 from GIA and ITRF and assuming that they are not correlated yields an uncertainty on the GMSL trend of ± 0.12 mm/yr over 1993-2017 ($1-\sigma$ level). In addition to the GIA correction and the TOPEX correction, we apply an elastic correction to the GMSL record of +0.10 mm/yr to account for the elastic deformations of the ocean bottom in response to modern melt of land ice (Frederikse et al., 2017; Lickley et al., 2018). The uncertainty in this correction
175 arises from the uncertainty associated to the computation of the elastic response of the solid Earth (mainly from the uncertainty associated to the procedure to solve the sea level equation, uncertainty in the choice of the Love numbers, uncertainty generated by the truncation degree of the spherical harmonics) and the uncertainty in the mass redistribution that cause the elastic deformation. Because the elastic response of the Earth is reasonably well defined (Mitrovica et al., 2011), the uncertainty in the elastic correction is largely dominated by the uncertainty in
180 the mass redistribution (Frederikse et al., 2017). The uncertainty on the mass redistribution is about $\pm 10\%$ on the current ice mass loss (e.g. [Blazquez et al., 2018](#)). It yields an uncertainty of $\pm 10\%$ on the elastic correction (because the elastic response of the Earth is linear). This uncertainty amounts ± 0.01 mm/yr which is very small. It is an order of magnitude smaller than the uncertainty considered in this study (see Table 1). So we neglect this source of uncertainty here.

185

The residual time correlated errors are separated into two different groups, depending on their correlation time scales. The first group gathers errors with short correlation time scales, i.e. lower than two months and between two months and one year. The second group gathers errors with long correlation time scales between 5 and 10 years. In the first group the errors are mainly due to the geophysical corrections (. ocean tides, atmospheric
190 corrections), to the altimeter corrections (sea-state bias correction, altimeter ionospheric corrections), to the orbital calculation, and to the potential altimeter instabilities (altimeter range and sigma-0 instabilities). At time scales below one year, the variability of the corrections' time series is dominated by errors, such that the variance of the error in each correction is estimated by the variance of the correction's time series. For errors with correlation time scales lower than 2 months, we estimated the standard deviation (σ) of the error from the correction's time series
195 filtered with a 2-month high-pass filter. Since the standard deviation of the errors depends on the different altimeter missions, the standard deviation has been separately estimated for each altimeter mission. We find $\sigma = 1.7$ mm over the T/P period, $\sigma = 1.5$ mm over the Jason-1 period, and $\sigma = 1.2$ mm over the Jason-2/3 period. For errors

with correlation time scale between 2 months and 1 year, we used the same approach and filtered the correction time series with a pass-band filter. In this case we find $\sigma = 1.3$ mm over the T/P period, $\sigma = 1.2$ mm over the Jason-1 period, and $\sigma = 1.0$ mm over the Jason-2/3 period. Unsurprisingly, the highest errors are obtained for T/P, and the lowest ones for Jason-2/3. This is because of: 1) larger altimeter range instabilities in T/P (Ablain et al., 2012; Beckley et al., 2017), 2) the presence of a 59-day signal error in the altimeter range of T/P (Zawadzki et al., 2018), and 3) because of the deterioration in the performance of atmospheric corrections in the early years of the altimetry era (Legeais et al., 2014). Note that Jason-1 shows also higher errors than Jason-2 and Jason-3 at time scales below 1 year (Couhert et al., 2015).

In the second group of residual time correlated errors, errors are due to the on-board microwave radiometer calibration, yielding instabilities in the wet troposphere correction, and also to the orbital calculation (Couhert et al., 2015). Since these errors are correlated at time scales longer than 5 years, they can not be estimated with the standard deviation of the correction time series, too short (25-year long) to sample the time correlation. For this group of residual-time correlated errors, we used simple models to represent the time correlation of the errors. For the wet troposphere correction, several studies Ablain, 2017 have identified long-term differences among the computed corrections from the different microwave radiometers and from the different atmospheric reanalyses (Dee et al., 2011). These studies report a difference in the wet tropospheric correction for GMSL in the range of ± 0.2 - 0.3 mm/yr for periods of 5 to 10 years. Here, we adopt a conservative approach and we model the error in wet tropospheric correction with a correlated error at 5 years with a standard deviation of 1.2 mm (1 σ level). The correlation is modelled with a Gaussian attenuation based on the wavelength of the errors: $e^{-\frac{1}{2}(\frac{t}{\lambda})^2}$ with $\lambda = 5$ years. In terms of trends, this residual time correlated error generates an uncertainty of ± 0.2 mm/yr over 5-year periods. For the error in the orbit calculation, comparisons of different orbit solutions showed differences of ± 0.05 mm/yr on 10 year time scales due to errors in the modelling of the Earth time varying gravity field (Couhert et al., 2015). We model this error with a correlated error at 10-year time scale with a standard deviation of 0.5 mm (1- σ level). The correlation is modelled by the same Gaussian distribution as before with $\lambda = 10$ years. In terms of trends, it corresponds to an uncertainty of ± 0.05 mm/yr over 10-year periods.

In the next section these different terms of the GMSL error budget are combined together to build the error variance-covariance matrix. Note that the different terms of the altimeter GMSL error budget described here are based on the current knowledge of altimetry measurement errors. As the altimetry record increases in length with new altimeter missions, the knowledge of the altimetry measurement also increases, and the description of the errors improves. This implies that the error variance-covariance matrix is expected to improve and change in the future.

3 The GMSL error variance-covariance matrix

In this section we derived the error variance-covariance matrix (Σ) of the GMSL from the error budget described in section 2. We assumed that all error sources shown in Table 1 are independent one to each other. Thus the Σ matrix is the sum of the individual variance-covariance matrix of each error source Σ_i in the error budget (see Figure 2). Each Σ_i matrix is calculated from a large number of random draws (> 1000) of simulated error signal using the model described in section 2 (either a bias, drift or time correlated signal) fed with a standard normal distribution. The resulting shape of each individual Σ_i matrix depends on the type of error (bias, drift or time correlated signal, see Figure 2). For the bias, the Σ_i matrix takes the shape of constant square blocks each side of the time occurrence of the bias correction (see for example the square matrix for TOPEX-A and TOPEX-B on the low left corner of Figure 2 along the diagonal). This square block shape means that the error in the bias correction generates an error on the GMSL which is fully correlated along time before and after the bias correction time, but which is not correlated along time for dates that are apart from the bias correction time. This is consistent with what we expect from a bias correction error. Note that in this article (and in climate change studies in general) we are interested only in GMSL changes, trends or acceleration but not in the mean time GMSL (which is the absolute reference of GMSL). Thus, we have removed from the GMSL time series the temporal mean over 1993-2017. The reference of the GMSL is thus arbitrary and assumed to be perfectly known. This is the reason because the reference of the GMSL is not affected by the biases correction error here.

For the drifts, the Σ_i matrix takes the shape of a horse saddle. This is because an error on the GMSL drift over a given period generates errors on the GMSL time series which are correlated when they are close in time and anti-correlated when they are on opposite side of the drift period.

For residual time correlated errors, the Σ_i matrix take the shape of a diagonal matrix with off diagonal terms of smaller amplitude. The further from the diagonal the off-diagonal terms are, the more attenuated they are. The attenuation rate is a Gaussian attenuation based on the wavelength of the time correlated errors ($e^{-\frac{1}{2}(\frac{t}{\lambda})^2}$), with various time-scales λ .

All individual Σ_i matrix are summed up together to build the total error variance-covariance matrix Σ of the altimetry-derived continuous GMSL record over 1993-2018 (see Figure 2). As expected, the dominant terms of the matrix are on the diagonal. They are largely due to the different sources of errors with correlation time scales below 1 year (first group of errors in section 2). The diagonal terms are the highest at the beginning of the altimetry period when T/P was at work. This is because of larger altimeter range instabilities in T/P, the presence of a 59-day signal error on the altimeter range of T/P and poorer performance of atmospheric corrections in the early years of the altimetry era (Legeais et al., 2014). The dominant off-diagonal terms are also found during the T/P period (in the lower left corner of the matrix, see Figure 2). The terms are induced by the TOPEX-A trend error and the large bias correction

uncertainty between TOPEX-A and TOPEX-B (because of the absence of inter-calibration phase between TOPEX-A and TOPEX-B).

4 GMSL uncertainty envelope

265 We estimated the GMSL uncertainty envelope from the square root of the diagonal terms of Σ (see Figure 3). As
expected, the GMSL time series shows a larger uncertainty during the T/P period (5 mm to 8 mm) than during the
Jason period (close to 4 mm). The bias correction uncertainty between TOPEX-A and TOPEX-B in February 1999
is also clearly visible with a 1-mm drop in the uncertainty after the switch to TOPEX side-B. Note that the uncertainty
envelope has a parabolic shape and shows smaller uncertainties during the beginning of the Jason-2 period (3.5
270 mm around 2008) than over the Jason-3 period (4.5 mm). This is not because Jason-1 and Jason-2 errors are
smaller than Jason-3's errors. Actually Jason 2 and Jason-3 errors are slightly smaller than Jason-1 errors thanks
to better orbit determination. The uncertainties are smaller during the Jason-1 and Jason-2 period because this
period is in the centre of the record. It benefits from prior and posterior data that covariate and help in reducing the
uncertainty when they are combined together. In contrast, the Jason-3 period is located at the end of the record
275 and does not benefit from posterior data to help reduce the uncertainty.

In Figure 4 we superimposed the GMSL time series (average of the GMSL time series in Figure 1) and the
associated uncertainty envelop. For the TOPEX-A period we tested three different curves with three different
corrections based on the removal of the Cal-1 mode (Beckley et al., 2017), on the comparison with tide gauges (
280 Watson et al., 2015; Ablain, 2017), or based on a sea level closure budget approach (Dieng et al., 2017). The
uncertainty envelop is centred on the corrected record for TOPEX-A drift with the correction based on Ablain et al.
(2017). As it has been expected, all the empirically corrected GMSL records are within the uncertainty envelop.

5 Uncertainty in GMSL trend and acceleration

285

The variance-covariance matrix can be used to derive the uncertainty on any metric based on the GMSL time
series. In this section we used the error variance-covariance matrix to estimate the uncertainty on the GMSL trend
and acceleration over any period of 5 years and more within 1993-2017.

290 Recently, several studies (Watson et al., 2015; Dieng et al., 2017; Nerem et al., 2018; WCRP Global Sea Level
Budget Group, 2018) have found a significant acceleration in the GMSL record from satellite altimetry (after

correction for the TOPEX-A drift). The occurrence of an acceleration in the record should not change the estimation of the trend when calculated with a least squared approach. However, it can affect the estimation of the uncertainty on the trend. To cope with this issue, we address here at the same time both the estimation of the trend and acceleration in the GMSL record. In order to obtain this objective, we used a second order polynomial as a predictor.

295 Considering the GMSL record has n observations, let X be an $n \times 3$ predictor where the first column contains only ones (representing the constant term), the second column contains the time vector (representing the linear term) and the third column contains the square of the time vector (representing the squared term). Let y be an $n \times 1$ vector of independent observations of the GMSL. Let ϵ be an $n \times 1$ vector of disturbances (GMSL non-linear and non-quadratic signals) and errors. Let β be the 3×1 vector of unknown parameters that we want to estimate,
 300 namely the GMSL y -intercept, the GMSL trend and the GMSL acceleration. Our linear regression model for the estimation of the GMSL trend and acceleration will thus be

$$y = X\beta + \epsilon$$

with

$$\epsilon \sim N(0, \Sigma)$$

305 where Σ is the variance-covariance matrix of the observation errors (estimated in the previous section). Σ is different from the identity because of the correlated noise (see section 2).

The most common method to estimate the GMSL trend and acceleration is the Ordinary Least Squares (OLS) estimator in its classical form (Cazenave and Llovel, 2010; Masters et al., 2012; Dieng et al., 2015; Nerem et al., 2018). This is also the most common method for estimating trends and accelerations in other climate essential
 310 variables (Hartmann, et al., 2014 and references therein). For these reasons, we turn here to the OLS to fit the linear regression model. The estimator of β with the OLS approach, noted $\hat{\beta}$ is

$$\hat{\beta} \sim (X^t X)^{-1} X^t y$$

In most cases, ϵ follows a $N(0, \sigma^2 I)$ distribution, which implies that $\hat{\beta}$ follows a Normal Law

$$\hat{\beta} = N(\beta, \sigma^2 (X^t X)^{-1})$$

315

The issue with this common framework is that the uncertainty of the trend and acceleration estimates does not take into account the correlated errors of the GMSL observations.

To address this issue, we used a more general formalism to integrate the GMSL error in the trend uncertainty estimation, following Ablain et al. (2009), Ribes et al. (2016) and IPCC AR5 (Hartmann, et al., 2014, see in particular
 320 Box 2.2 and Supplementary Material). The OLS estimator is left unchanged (and is still unbiased), but its distribution is revised to account for Σ , leading to:

$$\hat{\beta} = N(\beta, (X^t X)^{-1} (X^t \Sigma X) (X^t X)^{-1})$$

Note that this estimate is known to be less accurate than the General Least Square estimate (GLS, which is the optimal estimator in the case where $\Sigma \neq I$) in terms of the mean square error, because its variance is larger. A generalized least square estimate would probably help in narrowing slightly the trend uncertainty but the difference would likely be small as the GMSL time series is almost linear in time. Important advantages of using OLS here are (i) OLS is consistent with previous estimators of GMSL trends as well as estimators of trends in other essential climate variable than GMSL (e.g. Hartmann, et al., 2014), and (ii) the OLS best-estimate does not depend on the estimated variance-covariance matrix Σ .

Based on the matrix Σ defined in the previous section, and the OLS solution proposed before, we now estimate the GMSL trend (mm/yr) and acceleration (mm/yr²) uncertainties for any time span included in the period 1993-2017.

Results are synthetically displayed in




Figure 5 for trends and in Figure 6 for accelerations. On Figure 5, the top of the triangle indicates that the GMSL trend uncertainty over 1993-2017 is ± 0.4 mm/yr (CL 90%) and that the GMSL acceleration uncertainty over the same period is ± 0.07 mm/yr² (CL 90%, Figure 6). The GMSL acceleration uncertainty estimate is consistent with results of Watson et al., 2015, on the January 1993 to June 2014 time period where they find an uncertainty of

340 $\pm 0.058 \text{ mm} \cdot \text{yr}^{-2}$ at 1σ which corresponds to $\pm 0.096 \text{ mm/yr}^2$ at the 90% confidence level. This is slightly larger than the Nerem et al. (2018) estimate which is $\pm 0.025 \text{ mm/yr}^2$ at $1-\sigma$ level on the full 25-year altimetry era which corresponds to $\pm 0.041 \text{ mm/yr}^2$ at 90% confidence level. But the Nerem et al. (2018) estimate is likely underestimated as they only consider omission errors. The GMSL acceleration uncertainties have been calculated for all periods of 10 years and more within 1993-2017 (Figure 6). As expected, uncertainties tend to increase when
345 the period length decreases. At 10 years, the GMSL acceleration uncertainties are ranging from $\pm 0.3 \text{ mm/yr}^2$ over the T/P period to $\pm 0.25 \text{ mm/yr}^2$ over the Jason period. At 20 years they range between ± 0.12 and $\pm 0.08 \text{ mm/yr}^2$.

A cross-sectional analysis of the 10-year horizontal line on Figure 5 shows that the GMSL trend uncertainties over 10 years periods decreased from 1.0 mm/yr over the first decade to 0.5 mm/yr over the last one. The larger
350 uncertainty over the first decade is mainly due to the TOPEX-A drift error, but also to the large intermission bias uncertainty between TOPEX-A and TOPEX-B and, to a lesser extent, to the improvement of GMSL accuracy with Jason-2 and Jason-3. Note that the current GCOS requirement of 0.3 mm/yr uncertainty over 10 years (GCOS, 2011) is not met at the 90% confidence level. But the recent record over the last decade based on the Jason series is close to meet the GCOS requirement with a 90% CL.

355 Figure 5 can also be analysed by following the sides of the triangle. The results of this analysis are plotted on Figure 7. The plain line corresponds to the left side, read from bottom left to the top of the triangle. The dashed line corresponds to the right side, read from bottom right to the top of the triangle. As expected, both curves show a reduction of the trend uncertainty as the period over which trends are computed increases from 2 to 25 years. The difference between the two lines shows the reduction of GMSL errors thanks to the improvement of the
360 measurement in the latest altimetry missions. The lowest trend uncertainty is obtained with the last 20 years of the GMSL record: 0.35 mm/yr.

The periods for which the acceleration in sea level is significant at the 90% confidence level are shown in Figure 8. The acceleration is visible at the end of the record for periods of 10 years and longer. The GMSL acceleration is
365 0.12 mm/yr^2 with an uncertainty of 0.07 mm/yr^2 at 90% confidence level over the 25-year altimetry era. This proves that the acceleration observed in the GMSL evolution is statistically significant. It is worth noting that the different empirical TOPEX-A corrections yield very similar results (0.126 mm/yr^2 (Ablain, 2017) ; 0.120 mm/yr^2 (Dieng et al., 2017; Watson et al., 2015), 0.114 mm/yr^2 (Beckley et al., 2017)). This acceleration at the end of the record is due to an acceleration in the contribution to sea level from Greenland and from other contributions but to a lesser extent
370 (Chen et al., 2017; Dieng et al., 2017; Nerem et al., 2018). A small acceleration is also visible during the 1993-2005 period at the beginning of the record. This acceleration is likely due to the recovery from the Mount Pinatubo eruption in 1991 (Fasullo et al., 2016). Indeed, Church et al., 2005 showed that the impact of large volcanic eruptions on global ocean heat content is characterized by a rapid reduction in global ocean heat content during

the year following the eruption followed by a period of recovery of a few years when global ocean heat content
375 increases faster than before the eruption (see also (Gregory et al., 2006) and Delworth et al., 2005). The sea level
record starts in October 1992 which is 1.5 years after the eruption of Mount Pinatubo (15th of June 1991). At that
time the global ocean heat content was starting to recover with an increasing rate of rise (see Fasullo et al., 2016,
their fig.2) leading to an acceleration in sea level.

380 The period for which the trend in sea level is significant at the 90% confidence level is shown in





Figure 9. In periods when the acceleration is not significant, the second order polynomial that we used as a predictor
to estimate the trend and the acceleration does not hold anymore in principle. For these periods, we should turn
385 out a first order polynomial. The use of a first order polynomial does not affect the trend estimates, but only the
trend uncertainty estimates. We checked for differences in trend uncertainty when using either second order or first

order polynomial predictors. We found that these differences are negligible (not shown).



390 Figure 9 indicates that for periods of 5 years and longer, the trend in GMSL is always significant at 90% CL over the whole record. At the end of the record the trend tends to increase. This is consistent with the acceleration plot in Figure 6. Over the 25 years of satellite altimetry, we find a sea level rise of $.3.35 \pm 0.4$ mm/yr (90% CL), after correcting for the TOPEX-A GMSL drift. The differences due to the different TOPEX-A corrections are negligible (<0.05 mm.yr⁻¹).

395 **6 Conclusions**

In this study we have estimated the full GMSL error variance-covariance matrix over the satellite altimetry period. The matrix is available online (see section data). It provides to users a comprehensive description of the GMSL errors over the altimetry period. This matrix is based on the current knowledge of altimetry measurement errors. As the altimetry record increases in length with new altimeter missions, the knowledge of the altimetry measurement also increases and the description of the errors improves. Consequently, the error variance-

400

covariance matrix is expected to change and improve in the future – hopefully with a reduction of measurement uncertainty in new products.

405 The uncertainty of the GMSL here computed shows the reliability of altimetry measurements in order to accurately describe the evolution of the GMSL on all time scales from 10 days to 25 years. It also shows the reliability of altimetry measurements in order to estimate the trends and accelerations of the sea level. Along the altimetry record, we find that the uncertainty in each individual GMSL measurement decreases with time. It is smaller during the Jason era (2002-2018) than during the T/P period (1993-2002). Over the entire altimetry record, 1993-2017, we estimate the GMSL trend to 3.35 ± 0.4 mm/yr (90% CL, after correcting the TOPEX-A GMSL drift). We detect
410 also a significant GMSL acceleration over the 25-year period at 0.12 ± 0.07 mm/yr² (90% CL).

In this study, several assumptions have been made, that could be improved in the future. Firstly, the modelling of altimeter errors should be regularly revisited and improved to consider a better knowledge of errors (e.g. stability of wet troposphere corrections) and to consider future altimeter missions (e.g. Sentinel-3 and Sentinel-6 missions).
415 Concealing the mathematical formalism, OLS method has been applied because it is the most common approach used in the climate community to calculate trends in any climate data records. However this is not the optimal linear estimator. The use of a Generalized Least Square approach should involve some narrowing of trend or acceleration uncertainty. Another topic of concerns is the consideration of the internal and forced variability of the GMSL. Here we only considered the uncertainty in the GMSL due to the satellite altimeter instrument. In a future study, it would
420 be interesting to consider the partition of the GMSL into the forced response to anthropogenic forcing and the natural response to natural forcing and to the internal variability. Estimating the natural GMSL variability (e.g. using models) and considering it as an additional residual time correlated error, would allow to calculate the GMSL trend and acceleration representing the long-term evolution of GMSL in relationship with anthropogenic climate change.

Data

425 The global mean sea level error variance-covariance matrix is available online at <https://doi.org/10.17882/58344> (Ablain et al., 2018).

Acknowledgment

GS is funded by a FFABR (Finanziamento delle Attivita` Base di Ricerca) grant of the MIUR (Ministero dell'Istruzione, dell'Universita` e della Ricerca) and by a DiSPeA (Dipartimento di Scienze Pure e Applicate of the
430 Urbino University) grant.

This work was carried out as part of the Sea Level CCI (SL_cci) project (Climate Change Initiative programme) supported by ESA and the SALP (Service d'Altimétrie et de Localisation Précise) project supported by CNES for several years. We would also like to thank all contributors to these two projects, with special recognition to Thierry Guinle, SALP project manager at CNES.

435

References

- Ablain, M.: The TOPEX-A Drift and Impacts on GMSL Time Series, https://meetings.aviso.altimetry.fr/fileadmin/user_upload/tx_ausyclsseminar/files/Poster_OSTST17_GMSL_Drift_TOPEX-A.pdf, Miami, US (October, 2017). [online] Available from:
440 https://meetings.aviso.altimetry.fr/fileadmin/user_upload/tx_ausyclsseminar/files/Poster_OSTST17_GMSL_Drift_TOPEX-A.pdf (Accessed 8 November 2018), 2017.
- Ablain, M., Cazenave, A., Valladeau, G. and Guinehut, S.: A new assessment of the error budget of global mean sea level rate estimated by satellite altimetry over 1993–2008, *Ocean Science*, 5(2), 193–201, doi:10.5194/os-5-193-2009, 2009.
- Ablain, M., Philipps, S., Urvoy, M., Tran, N. and Picot, N.: Detection of Long-Term Instabilities on Altimeter Backscatter
445 Coefficient Thanks to Wind Speed Data Comparisons from Altimeters and Models, *Marine Geodesy*, 35(suppl. 1), 258–275, doi:10.1080/01490419.2012.718675, 2012.
- Ablain, M., Cazenave, A., Larnicol, G., Balmaseda, M., Cipollini, P., Faugère, Y., Fernandes, M. J., Henry, O., Johannessen, J. A., Knudsen, P., Andersen, O., Legeais, J., Meyssignac, B., Picot, N., Roca, M., Rudenko, S., Scharffenberg, M. G., Stammer, D., Timms, G. and Benveniste, J.: Improved sea level record over the satellite altimetry era (1993–2010) from the
450 Climate Change Initiative project, *Ocean Science*, 11(1), 67–82, doi:10.5194/os-11-67-2015, 2015.
- Ablain, M., Legeais, J. F., Prandi, P., Marcos, M., Fenoglio-Marc, L., Dieng, H. B., Benveniste, J. and Cazenave, A.: Satellite Altimetry-Based Sea Level at Global and Regional Scales, *Surveys in Geophysics*, 38(1), 7–31, doi:10.1007/s10712-016-9389-8, 2017.
- Ablain, M., Meyssignac, B., Zawadzki, L., Jugier, R., Ribes, A., Cazenave, A. and Picot, N.: Error variance-covariance matrix
455 of global mean sea level estimated from satellite altimetry (TOPEX, Jason 1, Jason 2, Jason 3), , doi:10.17882/58344, 2018.
- Beckley, B. D., Callahan, P. S., Hancock, D. W., Mitchum, G. T. and Ray, R. D.: On the “Cal-Mode” Correction to TOPEX Satellite Altimetry and Its Effect on the Global Mean Sea Level Time Series, *Journal of Geophysical Research: Oceans*, 122(11), 8371–8384, doi:10.1002/2017JC013090, 2017.
- Blazquez, A., Meyssignac, B., Lemoine, J., Berthier, E., Ribes, A. and Cazenave, A.: Exploring the uncertainty in GRACE
460 estimates of the mass redistributions at the Earth surface: implications for the global water and sea level budgets, *Geophysical Journal International*, 215(1), 415–430, doi:10.1093/gji/ggy293, 2018.
- Cazenave, A. and Llovel, W.: Contemporary Sea Level Rise, *Annual Review of Marine Science*, 2(1), 145–173, doi:10.1146/annurev-marine-120308-081105, 2010.
- Chen, X., Zhang, X., Church, J. A., Watson, C. S., King, M. A., Monselesan, D., Legresy, B. and Harig, C.: The increasing
465 rate of global mean sea-level rise during 1993–2014, *Nature Climate Change*, 7(7), 492–495, doi:10.1038/nclimate3325, 2017.

- Church, J. A., White, N. J. and Arblaster, J. M.: Significant decadal-scale impact of volcanic eruptions on sea level and ocean heat content, *Nature*, 438(7064), 74–77, doi:10.1038/nature04237, 2005.
- 470 Couhert, A., Cerri, L., Legeais, J.-F., Ablain, M., Zelensky, N. P., Haines, B. J., Lemoine, F. G., Bertiger, W. I., Desai, S. D. and Otten, M.: Towards the 1mm/y stability of the radial orbit error at regional scales, *Advances in Space Research*, 55(1), 2–23, doi:10.1016/j.asr.2014.06.041, 2015.
- 475 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N. and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553–597, doi:10.1002/qj.828, 2011.
- Delworth, T. L., Ramaswamy, V. and Stenchikov, G. L.: The impact of aerosols on simulated ocean temperature and heat content in the 20th century, *Geophys. Res. Lett.*, 32(24), L24709, doi:10.1029/2005GL024457, 2005.
- 480 Dieng, H. B., Cazenave, A., von Schuckmann, K., Ablain, M. and Meyssignac, B.: Sea level budget over 2005–2013: missing contributions and data errors, *Ocean Sci.*, 11(5), 789–802, doi:10.5194/os-11-789-2015, 2015.
- Dieng, H. B., Cazenave, A., Meyssignac, B. and Ablain, M.: New estimate of the current rate of sea level rise from a sea level budget approach, *Geophysical Research Letters*, 44(8), 3744–3751, doi:10.1002/2017GL073308, 2017.
- Fasullo, J. T., Nerem, R. S. and Hamlington, B.: Is the detection of accelerated sea level rise imminent?, *Scientific Reports*, 6(1), doi:10.1038/srep31245, 2016.
- 485 Frederikse, T., Riva, R. E. M. and King, M. A.: Ocean Bottom Deformation Due To Present-Day Mass Redistribution and Its Impact on Sea Level Observations: OCEAN BOTTOM DEFORMATION AND SEA LEVEL, *Geophys. Res. Lett.*, 44(24), 12,306–12,314, doi:10.1002/2017GL075419, 2017.
- 490 GCOS: Systematic Observation Requirements for Satellite-Based Data Products for Climate (2011 Update) – Supplemental details to the satellite-based component of the “Implementation Plan for the Global Observing System for Climate in Support of the UNFCCC (2010 Update), WMO., 2011.
- Gregory, J. M., Lowe, J. A. and Tett, S. F. B.: Simulated Global-Mean Sea Level Changes over the Last Half-Millennium, *J. Climate*, 19(18), 4576–4591, doi:10.1175/JCLI3881.1, 2006.
- 495 Hartmann, D. L., Klein Tank, A. M. G., Rusticucci, M., Alexander, L. V., Brönnimann, S., Charabi, Y. A. R. and Zhai, P.: Observations: Atmosphere and Surface, in *Climate Change 2013 - The Physical Science Basis : Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 159–254, Cambridge University Press, Cambridge., 2014.
- Henry, O., Ablain, M., Meyssignac, B., Cazenave, A., Masters, D., Nerem, S. and Garric, G.: Effect of the processing methodology on satellite altimetry-based global mean sea level rise over the Jason-1 operating period, *Journal of Geodesy*, 88(4), 351–361, doi:10.1007/s00190-013-0687-3, 2014.
- 500 Legeais, J.-F., Ablain, M. and Thao, S.: Evaluation of wet troposphere path delays from atmospheric reanalyses and radiometers and their impact on the altimeter sea level, *Ocean Science*, 10(6), 893–905, doi:10.5194/os-10-893-2014, 2014.

- 505 Legeais, J.-F., Ablain, M., Zawadzki, L., Zuo, H., Johannessen, J. A., Scharffenberg, M. G., Fenoglio-Marc, L., Fernandes, M. J., Andersen, O. B., Rudenko, S., Cipollini, P., Quartly, G. D., Passaro, M., Cazenave, A. and Benveniste, J.: An improved and homogeneous altimeter sea level record from the ESA Climate Change Initiative, *Earth System Science Data*, 10(1), 281–301, doi:<https://doi.org/10.5194/essd-10-281-2018>, 2018.
- Lickley, M. J., Hay, C. C., Tamisiea, M. E. and Mitrovica, J. X.: Bias in Estimates of Global Mean Sea Level Change Inferred from Satellite Altimetry, *J. Climate*, 31(13), 5263–5271, doi:[10.1175/JCLI-D-18-0024.1](https://doi.org/10.1175/JCLI-D-18-0024.1), 2018.
- 510 Masters, D., Nerem, R. S., Choe, C., Leuliette, E., Beckley, B., White, N. and Ablain, M.: Comparison of Global Mean Sea Level Time Series from TOPEX/Poseidon, Jason-1, and Jason-2, *Marine Geodesy*, 35(suppl. 1), 20–41, doi:[10.1080/01490419.2012.717862](https://doi.org/10.1080/01490419.2012.717862), 2012.
- Melini, D. and Spada, G.: Some remarks on Glacial Isostatic Adjustment modelling uncertainties, *Geophysical Journal International*, 218(1), 401–413, doi:[10.1093/gji/ggz158](https://doi.org/10.1093/gji/ggz158), 2019.
- 515 Meyssignac, B., Boyer, T., Zhao, Z., Hakuba, M. Z., Landerer, F. W., Stammer, D., Kato, S., Köhl, A., Ablain, M., Abraham, J. P., Blazquez, A., Cazenave, A., Church, J. A., Rebecca, C., Cheng, L., Domingues, C. and Giglio, D.: Measuring Global Ocean Heat Content to estimate the Earth Energy Imbalance, In press, *Front. Mar. Sci. - Ocean Observation*, 2018.
- Mitrovica, J. X., Gomez, N., Morrow, E., Hay, C., Latychev, K. and Tamisiea, M. E.: On the robustness of predictions of sea level fingerprints: On predictions of sea-level fingerprints, *Geophysical Journal International*, 187(2), 729–742, doi:[10.1111/j.1365-246X.2011.05090.x](https://doi.org/10.1111/j.1365-246X.2011.05090.x), 2011.
- 520 Nerem, R. S., Beckley, B. D., Fasullo, J. T., Hamlington, B. D., Masters, D. and Mitchum, G. T.: Climate-change–driven accelerated sea-level rise detected in the altimeter era, *PNAS*, 201717312, doi:[10.1073/pnas.1717312115](https://doi.org/10.1073/pnas.1717312115), 2018.
- Quartly, G. D., Legeais, J.-F., Ablain, M., Zawadzki, L., Fernandes, M. J., Rudenko, S., Carrère, L., García, P. N., Cipollini, P., Andersen, O. B., Poisson, J.-C., Mbajon Njiche, S., Cazenave, A. and Benveniste, J.: A new phase in the production of quality-controlled sea level data, *Earth System Science Data*, 9(2), 557–572, doi:[10.5194/essd-9-557-2017](https://doi.org/10.5194/essd-9-557-2017), 2017.
- 525 Ribes, A., Corre, L., Gibelin, A.-L. and Dubuisson, B.: Issues in estimating observed change at the local scale - a case study: the recent warming over France: estimating observed warming at the local scale, *International Journal of Climatology*, 36(11), 3794–3806, doi:[10.1002/joc.4593](https://doi.org/10.1002/joc.4593), 2016.
- Rudenko, S., Neumayer, K.-H., Dettmering, D., Esselborn, S., Schone, T. and Raimondo, J.-C.: Improvements in Precise Orbits of Altimetry Satellites and Their Impact on Mean Sea Level Monitoring, *IEEE Transactions on Geoscience and Remote Sensing*, 55(6), 3382–3395, doi:[10.1109/TGRS.2017.2670061](https://doi.org/10.1109/TGRS.2017.2670061), 2017.
- 530 Slangen, A. B. A., Adloff, F., Jevrejeva, S., Leclercq, P. W., Marzeion, B., Wada, Y. and Winkelmann, R.: A Review of Recent Updates of Sea-Level Projections at Global and Regional Scales, *Surveys in Geophysics*, 38(1), 385–406, doi:[10.1007/s10712-016-9374-2](https://doi.org/10.1007/s10712-016-9374-2), 2017.
- Spada, G.: Glacial Isostatic Adjustment and Contemporary Sea Level Rise: An Overview, *Surveys in Geophysics*, 38(1), 153–185, doi:[10.1007/s10712-016-9379-x](https://doi.org/10.1007/s10712-016-9379-x), 2017.
- 535 Thao, S., Eymard, L., Obligis, E. and Picard, B.: Trend and Variability of the Atmospheric Water Vapor: A Mean Sea Level Issue, *Journal of Atmospheric and Oceanic Technology*, 31(9), 1881–1901, doi:[10.1175/JTECH-D-13-00157.1](https://doi.org/10.1175/JTECH-D-13-00157.1), 2014.

Valladeau, G., Legeais, J. F., Ablain, M., Guinehut, S. and Picot, N.: Comparing Altimetry with Tide Gauges and Argo Profiling Floats for Data Quality Assessment and Mean Sea Level Studies, *Marine Geodesy*, 35(suppl. 1), 42–60, doi:10.1080/01490419.2012.718226, 2012.


540 Watson, C. S., White, N. J., Church, J. A., King, M. A., Burgette, R. J. and Legresy, B.: Unabated global mean sea-level rise over the satellite altimeter era, *Nature Climate Change*, 5(6), 565–568, doi:10.1038/nclimate2635, 2015.

WCRP Global Sea Level Budget Group: Global sea-level budget 1993–present, *Earth System Science Data*, 10(3), 1551–1590, doi:10.5194/essd-10-1551-2018, 2018.

545 Zawadzki, L. and Ablain, M.: Accuracy of the mean sea level continuous record with future altimetric missions: Jason-3 vs. Sentinel-3a, *Ocean Sci.*, 12(1), 9–18, doi:10.5194/os-12-9-2016, 2016.


Zawadzki, L., Ablain, M., Carrere, L., Ray, R. D., Zelensky, N. P., Lyard, F., Guillot, A. and Picot, N.: Investigating the 59-Day Error Signal in the Mean Sea Level Derived From TOPEX/Poseidon, Jason-1, and Jason-2 Data With FES and GOT Ocean Tide Models, *IEEE Transactions on Geoscience and Remote Sensing*, 56(6), 3244–3255, doi:10.1109/TGRS.2018.2796630, 2018.

550




555 Figure 1: Evolution of GMSL time series (corrected for TOPEX-A drift using Ablain (2017) TOPEX-A correction) from six different groups (AVISO/CNES, CSIRO, University of Colorado, SL_cci/ESA, NASA/GSFC, NOAA) products. The SL_cci/ESA covers a period from January 1993 to December 2016 while all other products cover the full 25-year period (January 1993 to December 2017). Seasonal (annual and semi-annual) signals have been removed and a 6-month smoothing has been applied. An averaged solution has been computed from the six

560 groups. GMSL time series have the same average on the 1993-2015 period (common period) and the averaged solution starts at zero in 1993. The averaged solution without TOPEX-A correction has also been represented. A GIA correction of -0.3 mm/yr has been subtracted to each data set. A correction of +0.10 mm/yr due to the deformations of the ocean bottom in response to modern melt of land ice (**Frederikse et al., 2017; Lickley et al., 2018**) has also been added.




565 Figure 2: Error variance-covariance matrix of altimeter GMSL on the 25-year period (January 1993 to December 2017).




570 Figure 3: Evolution in time of GMSL measurement uncertainty within a 90 % confidence level (1.65σ) on the 25-year period (January 1993 to December 2017).

575



580 **Figure 4: Evolution of the AVISO GMSL with different TOPEX-A corrections. On the black, red and green curves, the TOPEX-A drift correction has been respectively applied based on Ablain (2017), Watson et al. (2015), Dieng et al. (2017) and Beckley et al. (2017). The uncertainty envelope, as well as the trend and acceleration uncertainties are given to a 90% confidence level (1.65σ). Seasonal (annual and semi-annual) signals removed and 6-month smoothing applied. A GIA correction of -0.3 mm/yr has been subtracted to each data set. A correction of $+0.10 \text{ mm/yr}$ due to the deformations of the ocean bottom in response to modern melt of land ice (Frederikse et al., 2017; Lickley et al., 2018) has also been added.**

585




590

Figure 5: GMSL trend uncertainties (mm/yr) estimated for all altimeter periods within the 25-year period (January 1993 to December 2017). The confidence level is 90 % (1.65σ). Each colored pixel represents respectively the half-size of the 90% confidence range in the GMSL trend. Values are given in mm/yr. The vertical axis indicates the length of the period (ranging from 1 to 25 years) considered in the computation of the trend, while the

595


horizontal axis indicates the centre date of the period (for example 2000 for the 20-year period 1990-2009).



600


Figure 6: GMSL acceleration uncertainties (mm/yr²) estimated for all the altimeter periods within the 25-year period (January 1993 to December 2017). The confidence level is 90 % (1.65σ). Each colored pixel represents respectively the half-size of a 90% confidence range in the GMSL acceleration. Values are given in mm/yr². The vertical axis indicates the length of the period (ranging from 1 to 25 years) considered in the computation of the acceleration while the horizontal axis indicates the centre date of the period (for example 2000 for the 20-year period 1990-2009).

605




610

Figure 7: Evolution of the GMST trend uncertainties within a 90% confidence level (1.65σ) versus the altimeter period length from January 1993 to December 2017 on plain curve and from December 2017 to January 1993 on the dashed curve.



620 Figure 8: GMSL acceleration using the AVISO GMSL time series corrected for the TOPEX-A drift using the
 correction proposed by Ablain (2017): the acceleration in the shaded areas is not significant (lower than the
 acceleration uncertainties at the 90% confidence level). The length of the window (in years) is represented on the
 vertical axis and the central date of the used window (in years) is represented on the horizontal axis.



630 **Figure 9: GMSL trends using the AVISO GMSL timeseries corrected for the TOPEX-A drift using the correction proposed by (Ablain, 2017). The length of the window (in years) is represented on the vertical axis and the central date of the window used (in years) is represented on the horizontal axis. A GIA correction of -0.3 mm/yr has been subtracted. A correction of $+0.10$ mm/yr due to the deformations of the ocean bottom in response to modern melt of land ice (Frederikse et al., 2017; Lickley et al., 2018) has also been added.**

635

Source of errors	Error category	Uncertainty level (at 1 σ)	References
High frequency errors: altimeter noise, geophysical corrections, orbits ...	Correlated errors ($\lambda = 2$ months)	$\sigma = 1.7$ mm for TOPEX period $\sigma = 1.5$ mm for Jason-1 period. $\sigma = 1.2$ mm for Jason-2/3 period.	Calculation explained in this paper
Medium frequency errors: geophysical corrections, orbits ..	Correlated errors ($\lambda = 1$ year)	$\sigma = 1.3$ mm for TOPEX period $\sigma = 1.2$ mm for Jason-1 period. $\sigma = 1$ mm for Jason-2/3 period.	Calculation explained in this paper
Large frequency errors: wet troposphere correction	Correlated errors ($\lambda = 5$ years)	$\sigma = 1.1$ mm over all the period (\Leftrightarrow to 0.2 mm/yr for 5 years)	(Legeais et al., 2014; Thao et al., 2014)
Large frequency errors: orbits (Gravity fields)	Correlated errors ($\lambda = 10$ years)	$\sigma = 1.12$ mm over TOPEX period (no GRACE data) $\sigma = 0.5$ mm over Jason period (\Leftrightarrow to 0.05 mm/yr for 10 years)	(Couhert et al., 2015; Rudenko et al., 2017)
Altimeter instabilities on TOPEX-A and TOPEX-B	Drift error	$\delta = 0.7$ mm/yr on TOPEX-A period $\delta = 0.1$ mm/yr on TOPEX-B period	(Ablain, 2017; Beckley et al., 2017; Watson et al., 2015)
Long-term drift errors: orbit (ITRF) and GIA	Drift error	$\delta = 0.12$ mm/yr over 1993-2017	(Couhert et al., 2015; Spada, 2017)
GMSL bias errors to link altimetry missions together	Bias errors	$\Delta = 2$ mm for TP-A/TP-B $\Delta = 0.5$ mm for TP-B/J1, J1/J2, J2/J3.	(Zawadzki et al., 2018)

640

Table 1: Altimetry GMSL error budget given at 1-sigma